

*This table is a portion of the CCC PEP Quick Guide. It is intended to be used in conjunction with the Quick Guide and not as a standalone document.*

### Testing Recommendations for the Exposed Person

Recommendations		Baseline testing	Initial follow-up	Final follow-up
PEPline 2017	HCV+ SP (source person) <sup>1</sup>  or SP has potential HCV risk factors <sup>1</sup>	HCV Ab <sup>2</sup>	6 weeks <sup>3</sup>  HCV RNA  (HCV viral load)	≥6 months <sup>4,5</sup>  HCV Ab <sup>2</sup>
	SP HCV status unknown <sup>1</sup>  or SP is known and has no known HCV risk factors <sup>1</sup>		Optional 6 week HCV RNA	
CDC 2016 <sup>6</sup> (No longer recommended or posted. Pending revised recommendations from CDC)	All source persons <sup>1</sup>	HCV Ab <sup>2</sup>	≥3 weeks HCV RNA	Optional: ≥6 month HCV Ab <sup>2</sup>
CDC 2001 <sup>7</sup>		HCV Ab and ALT	If earlier diagnosis is desired: HCV RNA at 4-6 weeks	4-6 months HCV Ab and ALT

Abbreviations: HCV+ = hepatitis C positive; SP = source person; Ab = antibody; ALT = aminotransferase

- For purposes of initial post-exposure management, a source person is considered HCV-positive if either HCV antibody or HCV RNA (HCV viral load) is positive. Positive HCV antibody, however does not always indicate infectivity because: some persons eradicate HCV naturally but retain HCV antibody; and some patients have been successfully treated but retain HCV antibody. Persons with active HCV infection can have fluctuating HCV RNA (viral load) as well as undetectable viral load (and are presumably un-infectious at that time when viral load was undetectable).
- If the HCV antibody is positive at any point, follow-up HCV RNA testing is required. Persons with confirmed positive HCV RNA results should be referred for further evaluation and care.
- CDC Guidelines recommend the initial HCV follow-up with HCV RNA ≥ 3 weeks or HCV Ab ≥ 6 months with confirmatory RNA for those who test positive. A PEPline-recommended alternative is to do initial testing at 6 weeks, which coincides with the first HIV follow-up test. There are no data that establish a

clinical advantage to testing at 3 weeks vs. 6 weeks [Glynn, et al, Busch, et al, Hajarizadeh, et al]. HCV RNA becomes detectable beginning at 3 weeks. Testing earlier than 6 weeks can be performed at the discretion of the managing clinician, especially if preliminary assessment is needed. Positive HCV RNA indicates likely infection. However, approximately 25% of new infections will clear spontaneously [Naggie, et al]. Refer to an experienced provider for additional counseling, testing, and follow-up if positive.

**4** In HCV infection, HCV RNA can be transiently undetectable [Mosley, et al]. In addition, HCV antibodies develop slowly. Therefore, even though an early initial negative HCV RNA can be preliminarily reassuring, the PELine recommends further HCV antibody testing at  $\geq 6$  months post-exposure to confirm transmission did not occur.

**5** An interval (i.e. 12-16 week) HCV antibody test may provide some reassurance for exposed persons in most instances (and align with HIV surveillance). However: (a) testing at this time point may not impact overall exposure management significantly, and (b) it is not sufficiently sensitive to exclude HCV transmission. Even at 15 weeks, only about 80% of HCV infected persons will have positive HCV Ab [MMWR rr5005a1]. Therefore, a  $\geq 6$ -month HCV antibody test is considered to be conclusive in excluding HCV acquisition:  $\geq 97\%$  will be positive at 6 months post-exposure [MMWR rr5005a1].

**6** Please note, no longer available on the CDC website, formerly “Updated Information for Healthcare Personnel Potentially Exposed to Hepatitis C Virus (HCV): Recommended Testing and Follow-up (CDC)”. Nov 2016.

**7** Updated U.S. Public Health Service Guidelines for the Management of Occupational Exposures to HBV, HCV, and HIV and Recommendations for Postexposure Prophylaxis. MMWR 2001; 50 (RR11): 1-42. Note regarding exposed persons with symptoms: Symptoms of a viral illness compatible with acute HCV at any point up to 6 months post-exposure should prompt immediate evaluation for HCV (HCV antibody, viral load and LFTs).

Note regarding availability and feasibility of HCV RNA testing: HCV RNA testing might not be available or feasible at all institutions. If it is not possible to obtain the recommended HCV RNA testing, surveillance using antibody testing is essential in assessing HCV transmission.

Note regarding hepatic enzyme testing: The PELine does not recommend routine liver enzyme testing for follow-up because of the possibility of abnormal results from causes other than HCV.

#### References cited

Glynn SA, Wright DJ, Kleinman SH, et al. Dynamics of viremia in early hepatitis C virus infection.

Transfusion. 2005 June; 45(6):994–1002. doi: 10.1111/j.1537-2995.2005.04390.x

Busch MP, Shafer KAP. Acute-phase hepatitis C virus infection: implications for research, diagnosis, and treatment. Clin Infect Dis. (2005) 40 (7):959-961. doi: 10.1086/428583

Hajarizadeh B, Grebely J, Applegate T, et al. Dynamics of HCV RNA levels during acute hepatitis C virus infection. J Med Virol. 2014 Oct; 86(10): 1722–1729. doi: 10.1002/jmv.24010

Naggie S, Holland DP, Sulkowski MS, et al. Hepatitis C virus postexposure prophylaxis in the healthcare worker: why direct-acting antivirals don't change a thing. Clin Infect Dis. 2017 Jan 1;64(1):92-99. Epub 2016 Sep 28. doi: 10.1093/cid/ciw656

Mosley JW, Operskalski EA, Tobler LH, et al. The course of hepatitis C viraemia in transfusion recipients prior to availability of antiviral therapy. J Viral Hepat. 2008 Feb;15(2):120-8. doi: 10.1111/j.1365-2893.2007.00900.x

Recommendations for preventing transmission of infections among chronic hemodialysis patients.

MMWR. April 27, 2001;50(RR05):1-43. <http://www.cdc.gov/mmwr/preview/mmwrhtml/rr5005a1.htm>